

Attacks on antidepressants: signs of deep-seated stigma?



Psychiatry is used to being attacked by external parties with antidiagnosis and antitreatment agendas. However, the recent disclosure that a doctor (Professor Peter Gøtzsche) had joined a new group, the Council for Evidence-based Psychiatry, whose launch was accompanied by newspaper headlines such as “Antidepressants do more harm than good, research says” and “Psychiatric drugs are doing us more harm than good” in *The Times* and *The Guardian* plumbs a new nadir in irrational polemic. What is especially worrying is that this doctor is a co-founder of the Nordic Cochrane collaboration, an initiative set up to provide the best evidence for clinical practitioners. What is the truth about antidepressant efficacy and adverse effects, and why would Professor Gøtzsche apparently suspend his training in evidence analysis for popular polemic?

Depression is a serious and recurrent disorder that is currently the largest cause of disability in Europe¹ and is projected to be the leading cause of morbidity in high-income countries by 2030.² Antidepressants have an impressive effect size in the treatment of acute cases of depression, with a number needed to treat of around six.³ For example, the recently updated Cochrane review of amitriptyline,⁴ which involved 18 randomised controlled trials and 1987 participants, shows that it is significantly more effective than placebo in achieving acute response (odds ratio 2.67, 95% CI 2.21–3.23), and that significantly fewer participants allocated to amitriptyline than to placebo withdrew from trials because of treatment inefficacy. How can this finding represent more harm than good? A smaller proportion of treated patients withdrew because of side-effects and the pattern of results was the same in industry-sponsored and independently funded trials.⁴ Indeed, in general, effect sizes for psychiatric indications do not differ from those of drugs used in physical medicine.⁵ Moreover, antidepressants have an impressive ability to prevent recurrence of depression, with a number needed to treat of around three, which makes them one of the most effective of all drugs.⁶

Suicide kills about 6000 people every year in the UK.⁷ Most of these people are depressed and more than 70% are not taking an antidepressant at the time of death.⁸ Blanket condemnation of antidepressants by lobby groups and colleagues risks increasing that proportion.

In countries where antidepressants are used properly, suicide rates have fallen substantially.⁹

Of course, all active drugs have adverse effects, but for the new antidepressants these are rarely severe or life-threatening, even in overdose situations. Indeed, the new antidepressants, especially the selective serotonin reuptake inhibitors, are some of the safest drugs ever made. In our experience, the vast majority of patients who choose to stay on them do so because they improve their mood and wellbeing rather than because they cannot cope with withdrawal symptoms when they stop. Many of the extreme examples of adverse effects given by the opponents of antidepressants are both rare and sometimes sufficiently bizarre as to warrant the description of an unexplained medical symptom. To attribute extremely unusual or severe experiences to drugs that appear largely innocuous in double-blind clinical trials is to prefer anecdote to evidence. The incentive of litigation might also distort the presentation of some of the claims.

Antipsychiatry groups usually claim that depressed patients should be treated with exercise and psychotherapy instead of drugs. However, little controlled evidence exists to support the use of psychotherapy as an alternative to antidepressants in major depression. Indeed, if psychotherapy had to be tested according to the same rules as drugs, then whether or not it could be licensed for this indication is questionable.¹⁰ Moreover, the implication that, unlike antidepressants, psychotherapy is free of adverse effects is highly misleading. Suicidal ideation¹¹ and even completed suicide¹² are recognised adverse effects with psychotherapy, and sexual interference with patients by therapists is a matter of concern.¹⁰ Finally, exercise treatment, as the recent Cochrane review concludes, “is moderately more effective than a control intervention for reducing symptoms of depression, but analysis of methodologically robust trials only shows a smaller effect” and exercise is no more acceptable to patients than are psychological or pharmacological treatments.¹³

What motivates doctors with a commitment to evidence-based practice to make such a series of flawed statements about antidepressants? We can only speculate. First, general practitioners (GPs) clearly see a lot of patients with minor somatic and

Lancet Psychiatry 2014

Published Online

May 27, 2014

[http://dx.doi.org/10.1016/S2215-0366\(14\)70232-9](http://dx.doi.org/10.1016/S2215-0366(14)70232-9)

For the article in *The Times* see <http://www.thetimes.co.uk/tto/health/news/article4076351.ece>

For the article in *The Guardian* see <http://www.theguardian.com/commentisfree/2014/apr/30/psychiatric-drugs-harm-than-good-ssri-antidepressants-benzodiazepines>

psychiatric problems. We know from our contacts with GP colleagues that such patients might not be who a GP with a conventional internal medicine background yearns to treat. It might be comforting to believe that treatment doesn't really matter. Second, contemporary bien pensant society remains resolutely dualist in its language and its understanding, and doctors are part of that society. The idea of a medicine for something lacking in substance (the mind) might seem a priori implausible, irrational, and undesirable. Third, the anti-psychiatry movement, although now long in the tooth, has revived itself with the recent conspiracy theory that the pharmaceutical industry, in league with psychiatrists, actively plots to create diseases and manufacture drugs no better than placebo. The anti-capitalist flavour of this belief resonates with anti-psychiatry's strong association with extreme or alternative political views.

Whatever the reasons, extreme assertions such as those made by Prof Gøtzsche are insulting to the discipline of psychiatry and at some level express and reinforce stigma against mental illnesses and the people who have them. The medical profession must challenge these poorly thought-out negative claims by one of its own very vigorously.

*David J Nutt, Guy M Goodwin, Dinesh Bhugra, Seena Fazel, Stephen Lawrie

Imperial College London, London, UK (DJN); University of Oxford, Oxford, UK (GMG, SF); Kings College London, London, UK (DB); and University of Edinburgh, Edinburgh, UK (SL)
d.nutt@imperial.ac.uk

DJN has received grants and personal fees from Lundbeck and GSK; and personal fees from Lilly, BMS, Otsuka, Servier, and Pfizer. GMG has received grants and personal fees from Servier and Lundbeck; personal fees from Teva, Otsuka, Takeda, Eli Lilly, Merck, GSK, and AstraZeneca; and grants from P1vital. SL has received research funding from Abbvie, Roche, and Pfizer in connection with genetic, brain imaging, and therapeutic studies of people with schizophrenia. He has also been paid by Janssen and Roche to speak at or chair educational meetings about schizophrenia, as well as to contribute to advisory boards about new antipsychotic treatments. The other authors declare no competing interests.

- 1 Wittchen HU, Jacobi F, Rehm J, et al. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol* 2011; **21**: 655–79.
- 2 Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006; **3**: e442.
- 3 Anderson IM, Ferrier IN, Baldwin RC, et al. Evidence-based guidelines for treating depressive disorders with antidepressants: a revision of the 2000 British Association for Psychopharmacology guidelines. *J Psychopharmacol* 2008; **22**: 343–96.
- 4 Leucht C, Huhn M, Leucht S. Amitriptyline versus placebo for major depressive disorder. *Cochrane Database Syst Rev* 2012; **12**: CD009138.
- 5 Leucht, S Hierl S, Kissling W, Dold M, Davis JM. Putting the efficacy of psychiatric and general medicine medication into perspective: review of meta-analyses. *Br J Psychiatry* 2012; **200**: 97–106.
- 6 Geddes J, Carney S, Davies C, et al. Relapse prevention with antidepressant drug treatment in depressive disorders: a systematic review. *Lancet* 2003; **361**: 653–61.
- 7 Office for National Statistics. Suicide rates in the United Kingdom, 2012 Registrations. <http://www.ons.gov.uk/ons/rel/subnational-health4/suicides-in-the-united-kingdom/2012/stb-uk-suicides-2012.html> (accessed May 20, 2014).
- 8 Fazel S, Grann M, Ahlner J, Goodwin G. Suicides by violent means in individuals taking SSRIs and other antidepressants: a post-mortem study in Sweden, 1992–2004. *J Clin Psychopharmacol* 2007; **27**: 503–6.
- 9 Isacson G, Holmgren A, Osby U, Ahlner J. Decrease in suicides among the individuals treated with antidepressants: a controlled study of antidepressants in suicide in Sweden 1995–2005. *Acta Psychiatr Scand* 2009; **120**: 37–44.
- 10 Nutt DJ, Sharpe M. Uncritical positive regard? Issues in the safety and efficacy of psychotherapy. *J Psychopharmacol* 2008; **22**: 3–6.
- 11 Bridge J A, Barbe R P, Birmaher B, et al. Emergent suicidality in a clinical psychotherapy trial for adolescent depression. *Am J Psychiatry* 2005; **162**: 2173–75.
- 12 Stone A. Suicide precipitated by psychotherapy. *Am J Psychotherapy* 1971; **25**: 18–28.
- 13 Cooney GM, Dwan K, Greig CA, Lawlor DA, Rimer J, Waugh FR, McMurdo M, Mead GE. Exercise for depression. *Cochrane Database Syst Rev* 2013; **9**: CD004366.